

q -poly-Bernoulli numbers and q -poly-Cauchy numbers with a parameter by Jackson's integrals

Takao Komatsu

School of Mathematics and Statistics
Wuhan University, Wuhan, 430072, China
komatsu@whu.edu.cn

MR Subject Classifications: 05A15, 05A19, 11B75, 33D45

Abstract

We define q -poly-Bernoulli polynomials $B_{n,\rho,q}^{(k)}(z)$ with a parameter ρ , q -poly-Cauchy polynomials of the first kind $c_{n,\rho,q}^{(k)}(z)$ and of the second kind $\hat{c}_{n,\rho,q}^{(k)}(z)$ with a parameter ρ by Jackson's integrals, which generalize the previously known numbers and polynomials, including poly-Bernoulli numbers $B_n^{(k)}$ and the poly-Cauchy numbers of the first kind $c_n^{(k)}$ and of the second kind $\hat{c}_n^{(k)}$. We investigate their properties connected with usual Stirling numbers and weighted Stirling numbers. We also give the relations between generalized poly-Bernoulli polynomials and two kinds of generalized poly-Cauchy polynomials.

1 Introduction

Let n and k be integers with $n \geq 0$, and let ρ be a real number parameter¹ with $\rho \neq 0$. Let q be a real number with $0 \leq q < 1$. Define q -poly-Bernoulli polynomials $B_{n,\rho,q}^{(k)}(z)$ with a parameter ρ by

$$\frac{\rho}{1 - e^{-\rho t}} \text{Li}_{k,q} \left(\frac{1 - e^{-\rho t}}{\rho} \right) e^{-tz} = \sum_{n=0}^{\infty} B_{n,\rho,q}^{(k)}(z) \frac{t^n}{n!}, \quad (1)$$

where $\text{Li}_{k,q}(z)$ is the q -polylogarithm function ([13]) defined by

$$\text{Li}_{k,q}(z) = \sum_{n=1}^{\infty} \frac{z^n}{[n]_q^k}.$$

Here,

$$[x]_q = \frac{1 - q^x}{1 - q}$$

¹ q was used as a parameter in [15, 5], but in this paper we use ρ in order to avoid confusions with q -integral.

is the q -number with $[0]_q = 0$ (see e.g. [1, (10.2.3)], [9]). Note that $\lim_{q \rightarrow 1} [x]_q = x$.

Notice that

$$\lim_{q \rightarrow 1} B_{n,\rho,q}^{(k)}(z) = B_{n,\rho}^{(k)}(z),$$

which is the poly-Bernoulli polynomial with a ρ parameter ([5]), and

$$\lim_{q \rightarrow 1} \text{Li}_{k,q}(z) = \text{Li}_k(z),$$

which is the ordinary polylogarithm function, defined by

$$\text{Li}_k(z) = \sum_{m=1}^{\infty} \frac{z^m}{m^k}. \quad (2)$$

In addition, when $z = 0$, $B_{n,\rho}^{(k)}(0) = B_{n,\rho}^{(k)}$ is the poly-Bernoulli number with a ρ parameter. When $z = 0$ and $\rho = 1$, $B_{n,1}^{(k)}(0) = B_n^{(k)}$ is the poly-Bernoulli number ([12]) defined by

$$\frac{\text{Li}_k(1 - e^{-t})}{1 - e^{-t}} = \sum_{n=0}^{\infty} B_n^{(k)} \frac{t^n}{n!}, \quad (3)$$

The poly-Bernoulli numbers are extended to the poly-Bernoulli polynomials ([2, 7]) and to the special multi-poly-Bernoulli numbers ([8]).

Let n and k be integers with $n \geq 0$, and let ρ be a real number parameter with $\rho \neq 0$. Let q be a real number with $0 \leq q < 1$. Define q -poly-Cauchy polynomials of the first kind $c_{n,\rho,q}^{(k)}(z)$ with a parameter ρ by

$$\begin{aligned} c_{n,\rho,q}^{(k)}(z) &= \rho^n \underbrace{\int_0^1 \cdots \int_0^1}_k \left(\frac{x_1 \cdots x_k - z}{\rho} \right)_n d_q x_1 \cdots d_q x_k \\ &= \rho^n n! \underbrace{\int_0^1 \cdots \int_0^1}_k \left(\frac{(x_1 \cdots x_k - z)/\rho}{n} \right) d_q x_1 \cdots d_q x_k, \end{aligned} \quad (4)$$

where $(x)_n = x(x-1) \cdots (x-n+1)$ ($n \geq 1$) with $(x)_0 = 1$.

Jackson's q -derivative with $0 < q < 1$ (see e.g. [1, (10.2.3)], [9]) is defined by

$$D_q f = \frac{d_q f}{d_q x} = \frac{f(x) - f(qx)}{(1-q)x}$$

and Jackson's q -integral ([1, (10.1.3)], [9]) is defined by

$$\int_0^x f(t) d_q t = (1-q)x \sum_{n=0}^{\infty} f(q^n x) q^n.$$

For example, when $f(x) = x^m$ for some nonnegative integer m ,

$$\begin{aligned} D_q f &= \frac{x^m - q^m x^m}{(1-q)x} \\ &= [m]_q x^{m-1} \end{aligned}$$

and

$$\begin{aligned} \int_0^x t^m d_q t &= (1-q)x \sum_{n=0}^{\infty} q^{mn} x^m q^n \\ &= (1-q)x^{m+1} \sum_{n=0}^{\infty} q^{n(m+1)} \\ &= \frac{x^{m+1}}{[m+1]_q}. \end{aligned}$$

Notice that

$$\lim_{q \rightarrow 1} c_{n,\rho,q}^{(k)}(z) = c_{n,\rho}^{(k)}(z),$$

which is the poly-Cauchy polynomial with a ρ parameter ([15]). In addition, when $z = 0$, $c_{n,\rho}^{(k)}(0) = c_{n,\rho}^{(k)}$ is the poly-Cauchy number with a ρ parameter ([15]). When $z = 0$ and $\rho = 1$, $c_{n,1}(0) = c_n^{(k)}$ is the poly-Cauchy number (of the first kind) ([14]), defined by the integral of the falling factorial:

$$\begin{aligned} c_n^{(k)} &= \underbrace{\int_0^1 \cdots \int_0^1}_{k} (x_1 \cdots x_k)_n dx_1 \cdots dx_k \\ &= n! \underbrace{\int_0^1 \cdots \int_0^1}_{k} \binom{x_1 \cdots x_k}{n} dx_1 \cdots dx_k. \end{aligned}$$

If $k = 1$, then $c_n^{(1)} = c_n$ is the classical Cauchy number ([6, 20]). The number $c_n/n!$ is sometimes referred to as the Bernoulli number of the second kind ([3, 10, 21]). The poly-Cauchy numbers of the first kind $c_n^{(k)}$ can be expressed in terms of the Stirling numbers of the first kind.

$$c_n^{(k)} = \sum_{m=0}^n \frac{(-1)^{n-m} S_1(n, m)}{(m+1)^k} \quad (n \geq 0, k \geq 1)$$

([14, Theorem 1]), where $S_1(n, m)$ is the (unsigned) Stirling number of the first kind, see [6], determined by the rising factorial:

$$x(x+1) \cdots (x+n-1) = \sum_{m=0}^n S_1(n, m) x^m. \quad (5)$$

The generating function of the poly-Cauchy numbers $c_n^{(k)}$ is given by

$$\text{Lif}_k(\ln(1+t)) = \sum_{n=0}^{\infty} c_n^{(k)} \frac{t^n}{n!} \quad (6)$$

([14, Theorem 2]), where $\text{Lif}_k(z)$ is called *polylogarithm factorial* function (or simply, *polyfactorial* function) defined by

$$\text{Lif}_k(z) = \sum_{m=0}^{\infty} \frac{z^m}{m!(m+1)^k}. \quad (7)$$

By this definition, k is not restricted to a positive integer in $c_n^{(k)}$. Similarly, define the poly-Cauchy numbers of the second kind $\hat{c}_n^{(k)}$ ([14]) by

$$\begin{aligned} \hat{c}_n^{(k)} &= \underbrace{\int_0^1 \cdots \int_0^1}_{k} (-x_1 \cdots x_k)_n dx_1 \cdots dx_k \\ &= n! \underbrace{\int_0^1 \cdots \int_0^1}_{k} \binom{-x_1 \cdots x_k}{n} dx_1 \cdots dx_k. \end{aligned}$$

If $k = 1$, then $\hat{c}_n^{(1)} = \hat{c}_n$ is the classical Cauchy number of the second kind ([6, 20]). The poly-Cauchy numbers of the second kind $\hat{c}_n^{(k)}$ can be expressed in terms of the Stirling numbers of the first kind.

$$\hat{c}_n^{(k)} = (-1)^n \sum_{m=0}^n \frac{S_1(n, m)}{(m+1)^k} \quad (n \geq 0, k \geq 1)$$

([14, Theorem 4]). The generating function of the poly-Cauchy numbers of the second kind $\hat{c}_n^{(k)}$ is given by

$$\text{Lif}_k(-\ln(1+t)) = \sum_{n=0}^{\infty} \hat{c}_n^{(k)} \frac{t^n}{n!} \quad (8)$$

([14, Theorem 5]).

The poly-Cauchy numbers have been considered as analogues of poly-Bernoulli numbers $B_n^{(k)}$. The poly-Cauchy numbers (of the both kinds) are extended to the poly-Cauchy polynomials ([11]), and to the poly-Cauchy numbers with a q parameter ([15]). The corresponding poly-Bernoulli numbers with a q parameter can be obtained in [5]. A different direction of generalizations of Cauchy numbers is about Hypergeometric Cauchy numbers ([17]). Arithmetical and combinatorial properties including sums of products have been studied ([16, 18, 19]).

In this paper, by using Jackson's q -integrals, as essential generalizations of the previously known numbers, including poly-Bernoulli numbers $B_n^{(k)}$, the poly-Cauchy numbers of the first kind $c_n^{(k)}$ and of the second kind $\hat{c}_n^{(k)}$, we introduce the concept about

q -analogues or extensions of the poly-Bernoulli polynomials $B_{n,\rho,q}^{(k)}(z)$ with a parameter, the poly-Cauchy polynomials of the first kind $c_{n,\rho,q}^{(k)}$ and of the second kind $\hat{c}_{n,\rho,q}^{(k)}$ with a parameter. We investigate their properties connected with usual Stirling numbers and weighted Stirling numbers. We also give the relations between generalize poly-Bernoulli polynomials and two kinds of generalized poly-Cauchy polynomials.

2 q -poly-Bernoulli polynomials with a parameter

Carlitz [4] defined the weighted Stirling numbers of the first kind $S_1(n, m, x)$ and of the second kind $S_2(n, m, x)$ by

$$\frac{(1-t)^{-x}(-\ln(1-t))^m}{m!} = \sum_{n=0}^{\infty} S_1(n, m, x) \frac{t^n}{n!} \quad (9)$$

and

$$\frac{e^{xt}(e^t - 1)^m}{m!} = \sum_{n=0}^{\infty} S_2(n, m, x) \frac{t^n}{n!}, \quad (10)$$

respectively. Note that Carlitz [4] used the notation $R_1(n, m, x)$ and $R(n, m, x)$ instead of $S_1(n, m, x)$ and $S_2(n, m, x)$, respectively. When $x = 0$, $S_1(n, m, 0) = S_1(n, m)$ and $S_2(n, m, 0) = S_2(n, m)$ are the (unsigned) Stirling number of the first kind and the Stirling number of the second kind, respectively.

The q -poly-Bernoulli polynomials with a parameter ρ can be expressed in terms of the weighted Stirling numbers of the second kind.

Theorem 1 *We have*

$$B_{n,\rho,q}^{(k)}(z) = \sum_{m=0}^n S_2\left(n, m, \frac{z}{\rho}\right) \frac{(-\rho)^{n-m} m!}{[m+1]_q^k}. \quad (11)$$

Proof. From (1), and using (10), we have

$$\begin{aligned} \sum_{n=0}^{\infty} B_{n,\rho,q}^{(k)}(z) \frac{t^n}{n!} &= \sum_{m=0}^{\infty} \frac{(-\rho)^{-m}}{[m+1]_q^k} e^{-\rho t(z/\rho)} (e^{-\rho t} - 1)^m \\ &= \sum_{m=0}^{\infty} \frac{(-\rho)^{-m} m!}{[m+1]_q^k} \sum_{n=m}^{\infty} S_2\left(n, m, \frac{z}{\rho}\right) \frac{(-\rho t)^n}{n!} \\ &= \sum_{n=0}^{\infty} \left(\sum_{m=0}^n S_2\left(n, m, \frac{z}{\rho}\right) \frac{(-\rho)^{n-m} m!}{[m+1]_q^k} \right) \frac{t^n}{n!}. \end{aligned}$$

Comparing the coefficients on both sides, we get the result. Notice that $S_2(n, m, x) = 0$ for $n < m$. ■

Corollary 1 For q -poly-Bernoulli numbers with a parameter ρ , we have

$$B_{n,\rho,q}^{(k)} = \sum_{m=0}^n S_2(n, m) \frac{(-\rho)^{n-m} m!}{[m+1]_q^k}. \quad (12)$$

3 q -poly-Cauchy polynomials of the first kind with a parameter

The q -poly-Cauchy polynomials of the first kind $c_{n,\rho,q}^{(k)}(z)$ with a parameter can be expressed in terms of the weighted Stirling numbers of the first kind $S_1(n, m, x)$. In this expression, k is not restricted to a positive integer.

Theorem 2 For integers n and k with $n \geq 0$, we have

$$\begin{aligned} c_{n,\rho,q}^{(k)}(z) &= \sum_{m=0}^n S_1(n, m) (-\rho)^{n-m} \sum_{i=0}^m \binom{m}{i} \frac{(-z)^i}{[m-i+1]_q^k} \\ &= \sum_{m=0}^n S_1\left(n, m, \frac{z}{\rho}\right) \frac{(-\rho)^{n-m}}{[m+1]_q^k}. \end{aligned} \quad (13)$$

Proof. From (4) and (5), we have

$$\begin{aligned} c_{n,\rho,q}^{(k)}(z) &= \rho^n \sum_{m=0}^n (-1)^{n-m} S_1(n, m) \underbrace{\int_0^1 \cdots \int_0^1}_{k} \left(\frac{x_1 \cdots x_k - z}{\rho} \right)^m d_q x_1 \cdots d_q x_k \\ &= \sum_{m=0}^n (-\rho)^{n-m} S_1(n, m) \sum_{i=0}^m \binom{m}{i} (-z)^{m-i} \underbrace{\int_0^1 \cdots \int_0^1}_{k} x_1^i \cdots x_k^i d_q x_1 \cdots d_q x_k \\ &= \sum_{m=0}^n (-\rho)^{n-m} S_1(n, m) \sum_{i=0}^m \binom{m}{i} \frac{(-z)^{m-i}}{[i+1]_q^k} \\ &= \sum_{m=0}^n (-\rho)^{n-m} S_1(n, m) \sum_{i=0}^m \binom{m}{i} \frac{(-z)^i}{[m-i+1]_q^k}. \end{aligned}$$

By using the relation, see [4, Eq. (5.2)],

$$S_1(n, m, x) = \sum_{i=0}^n \binom{m+i}{i} x^i S_1(n, m+i),$$

we obtain

$$\begin{aligned}
c_{n,\rho,q}^{(k)}(z) &= \sum_{i=0}^n \sum_{m=i}^n S_1(n, m) (-\rho)^{n-m} \binom{m}{i} \frac{(-z)^i}{[m-i+1]_q^k} \\
&= \sum_{i=0}^n \sum_{m=i}^{n+i} S_1(n, m) (-\rho)^{n-m} \binom{m}{i} \frac{(-z)^i}{[m-i+1]_q^k} \\
&= \sum_{i=0}^n \sum_{m=0}^n S_1(n, m+i) (-\rho)^{n-m-i} \binom{m+i}{i} \frac{(-z)^i}{[m+1]_q^k} \\
&= \sum_{m=0}^n \frac{(-\rho)^{n-m}}{[m+1]_q^k} \sum_{i=0}^m \binom{m+i}{i} \left(\frac{z}{\rho}\right)^i S_1(n, m+i) \\
&= \sum_{m=0}^n S_1\left(n, m, \frac{z}{\rho}\right) \frac{(-\rho)^{n-m}}{[m+1]_q^k}.
\end{aligned}$$

■

Corollary 2 *For q -poly-Cauchy numbers of the first kind with a parameter ρ , we have*

$$c_{n,\rho,q}^{(k)} = \sum_{m=0}^{\infty} S_1(n, m) \frac{(-\rho)^{n-m}}{[m+1]_q^k}. \quad (14)$$

Define the q -polyfactorial functions $\text{Lif}_{k,q}(z)$ by

$$\text{Lif}_{k,q}(z) = \sum_{n=0}^{\infty} \frac{z^n}{n! [n+1]_q^k}.$$

Notice that

$$\lim_{q \rightarrow 1} \text{Lif}_{k,q}(z) = \text{Lif}_k(z),$$

which is the ordinary polyfactorial function in (7). The generating function of $c_{n,\rho,q}^{(k)}$ is given by the following theorem.

Theorem 3 *We have*

$$\frac{1}{(1+\rho t)^{z/\rho}} \text{Lif}_{k,q}\left(\frac{\ln(1+\rho t)}{\rho}\right) = \sum_{n=0}^{\infty} c_{n,\rho,q}^{(k)}(z) \frac{t^n}{n!}. \quad (15)$$

Proof. By the first identity of Theorem 2

$$\begin{aligned}
\sum_{n=0}^{\infty} c_{n,\rho,q}^{(k)}(z) \frac{t^n}{n!} &= \sum_{n=0}^{\infty} \sum_{m=0}^n S_1(n, m) (-\rho)^{n-m} \sum_{i=0}^m \binom{m}{i} \frac{(-z)^i}{[m-i+1]_q^k} \frac{t^n}{n!} \\
&= \sum_{m=0}^{\infty} (-\rho)^{-m} \sum_{n=m}^{\infty} S_1(n, m) \frac{(-\rho t)^n}{n!} \sum_{i=0}^m \binom{m}{i} \frac{(-z)^i}{[m-i+1]_q^k} \\
&= \sum_{m=0}^{\infty} \frac{1}{m!} \left(\frac{\ln(1+\rho t)}{\rho} \right)^m \sum_{i=0}^m \binom{m}{i} \frac{(-z)^i}{[m-i+1]_q^k} \\
&= \sum_{i=0}^{\infty} \frac{(-z)^i}{i!} \sum_{m=i}^{\infty} \frac{1}{(m-i)! [m-i+1]_q^k} \left(\frac{\ln(1+\rho t)}{\rho} \right)^m \\
&= \sum_{n=0}^{\infty} \frac{(\ln(1+\rho t))^n}{n! \rho^n} \frac{1}{[n+1]_q^k} \sum_{i=0}^{\infty} \frac{1}{i!} \left(-\frac{z \ln(1+\rho t)}{\rho} \right)^i \\
&= \frac{1}{(1+\rho t)^{z/\rho}} \sum_{n=0}^{\infty} \frac{(\ln(1+\rho t))^n}{\rho^n} \frac{1}{n! [n+1]_q^k} \\
&= \frac{1}{(1+\rho t)^{z/\rho}} \text{Lif}_{k,q} \left(\frac{\ln(1+\rho t)}{\rho} \right).
\end{aligned}$$

■

4 q -poly-Cauchy polynomials of the second kind with a parameter

Let n and k be integers with $n \geq 0$ and $k \geq 1$, and let ρ be a real number parameter with $\rho \neq 0$. Define q -poly-Cauchy polynomials of the second kind $\widehat{c}_{n,\rho,q}^{(k)}(z)$ with a parameter ρ by

$$\begin{aligned}
\widehat{c}_{n,\rho,q}^{(k)}(z) &= \rho^n \underbrace{\int_0^1 \cdots \int_0^1}_k \left(\frac{-x_1 \cdots x_k + z}{\rho} \right)_n d_q x_1 \cdots d_q x_k \\
&= \rho^n n! \underbrace{\int_0^1 \cdots \int_0^1}_k \left(\frac{(-x_1 \cdots x_k + z)/\rho}{n} \right) d_q x_1 \cdots d_q x_k.
\end{aligned} \tag{16}$$

Notice that

$$\lim_{q \rightarrow 1} \widehat{c}_{n,\rho,q}^{(k)}(z) = \widehat{c}_{n,\rho}^{(k)}(z),$$

which is the poly-Cauchy polynomial of the second kind with a ρ parameter ([15]). In addition, when $z = 0$, $\widehat{c}_{n,\rho}^{(k)}(0) = \widehat{c}_{n,\rho}^{(k)}$ is the poly-Cauchy number of the second kind with

a ρ parameter ([15]). When $z = 0$ and $\rho = 1$, $\widehat{c}_{n,1}(0) = \widehat{c}_n^{(k)}$ is the poly-Cauchy number ([14]) given in (8).

The q -poly-Cauchy polynomials of the first kind $\widehat{c}_{n,\rho,q}^{(k)}(z)$ with a parameter can be expressed in terms of the weighted Stirling numbers of the first kind $S_1(n, m, x)$. In this expression, k is not restricted to positive integers.

Theorem 4 *For integers n and k with $n \geq 0$, we have*

$$\begin{aligned}\widehat{c}_{n,\rho,q}^{(k)}(z) &= (-1)^n \sum_{m=0}^n S_1(n, m) \rho^{n-m} \sum_{i=0}^m \binom{m}{i} \frac{(-z)^i}{[m-i+1]_q^k} \\ &= (-1)^n \sum_{m=0}^n S_1\left(n, m, -\frac{z}{\rho}\right) \frac{\rho^{n-m}}{[m+1]_q^k}.\end{aligned}\tag{17}$$

Proof. From (4) and (5), and using the relation [4, Eq. (5.2)], similarly to the proof of Theorem 2, we obtain the result. ■

Corollary 3 *For q -poly-Cauchy numbers of the second kind with a parameter ρ , we have*

$$\widehat{c}_{n,\rho,q}^{(k)} = (-1)^n \sum_{m=0}^{\infty} S_1(n, m) \frac{\rho^{n-m}}{[m+1]_q^k}.$$

Proof. Putting $z = 0$ in Theorem 4, we immediately get the result. ■

The generating function of $\widehat{c}_{n,\rho,q}^{(k)}$ is given by using the q -polyfactorial functions $\text{Lif}_{k,q}(z)$.

Theorem 5 *We have*

$$(1 + \rho t)^{z/\rho \text{Lif}_{k,q}} \left(-\frac{\ln(1 + \rho t)}{\rho} \right) = \sum_{n=0}^{\infty} \widehat{c}_{n,\rho,q}^{(k)}(z) \frac{t^n}{n!}.$$

Proof. Similarly to the proof of Theorem 3, by the first identity of Theorem 2, we obtain the result. ■

5 Several relations of q -poly-Bernoulli polynomials and q -poly-Cauchy polynomials

There exist orthogonality and inverse relations for weighted Stirling numbers ([4]). Namely, from the orthogonal relations

$$\sum_{l=m}^n (-1)^{n-l} S_2(n, l, x) S_1(l, m, x) = \sum_{l=m}^n (-1)^{l-m} S_1(n, l, x) S_2(l, m, x) = \delta_{m,n},$$

where $\delta_{m,n} = 1$ if $m = n$; $\delta_{m,n} = 0$ otherwise, we obtain the inverse relations

$$f_n = \sum_{m=0}^n (-1)^{n-m} S_1(n, m, x) g_m \iff g_n = \sum_{m=0}^n S_2(n, m, x) f_m. \quad (18)$$

Theorem 6 *For q -poly-Bernoulli and q -poly-Cauchy polynomials with a parameter, we have*

$$\sum_{m=0}^n S_1\left(n, m, \frac{z}{\rho}\right) \rho^{-m} B_{m,\rho,q}^{(k)}(z) = \frac{n!}{\rho^n [n+1]_q^k}, \quad (19)$$

$$\sum_{m=0}^n S_2\left(n, m, \frac{z}{\rho}\right) \rho^{-m} c_{m,\rho,q}^{(k)}(z) = \frac{1}{\rho^n [n+1]_q^k}, \quad (20)$$

$$\sum_{m=0}^n S_2\left(n, m, -\frac{z}{\rho}\right) \rho^{-m} \hat{c}_{m,\rho,q}^{(k)}(z) = \frac{(-1)^n}{\rho^n [n+1]_q^k}. \quad (21)$$

Remark. If $q \rightarrow 1$, then Theorem 6 is reduced to Theorem 3.2 in [5].

Proof. By Theorem 1, applying (18) with

$$f_m = \frac{m!}{(-\rho)^m [m+1]_q^k}, \quad g_n = \frac{B_{n,\rho,q}^{(k)}(z)}{(-\rho)^n}$$

and x is replaced by z/ρ , we get the identity (19). Similarly, by Theorem 2 and Theorem 4 we have the identities (20) and (21), respectively. $\blacksquare$

There are relations between two kinds of q -poly-Cauchy polynomials with a parameter.

Theorem 7 *For $n \geq 1$ we have*

$$(-1)^n \frac{c_{n,\rho,q}^{(k)}(z)}{n!} = \sum_{m=1}^n \binom{n-1}{m-1} \frac{\hat{c}_{m,\rho,q}^{(k)}(z)}{m!}, \quad (22)$$

$$(-1)^n \frac{\hat{c}_{n,\rho,q}^{(k)}(z)}{n!} = \sum_{m=1}^n \binom{n-1}{m-1} \frac{c_{m,\rho,q}^{(k)}(z)}{m!}. \quad (23)$$

Remark. Since $c_{n,1,q}^{(k)}(z) \rightarrow c_n^{(k)}(z)$ and $\hat{c}_{n,1,q}^{(k)}(z) \rightarrow \hat{c}_n^{(k)}(z)$ as $q \rightarrow 1$, Theorem 7 is reduced to Theorem 4.2 in [11].

Proof. We shall prove identity (23). The identity (22) can be proven similarly. By the definition of $\widehat{c}_{n,q}^{(k)}(z)$

$$\begin{aligned} (-1)^n \frac{\widehat{c}_{n,\rho,q}^{(k)}(z)}{n!} &= (-\rho)^n \underbrace{\int_0^1 \cdots \int_0^1}_k \binom{(-x_1 \cdots x_k + z)/\rho}{n} d_q x_1 \cdots d_q x_k \\ &= \rho^n \underbrace{\int_0^1 \cdots \int_0^1}_k \binom{(x_1 \cdots x_k - z)/\rho + (n-1)}{n} d_q x_1 \cdots d_q x_k. \end{aligned}$$

We use the well-known identity, see [22, p. 8],

$$\binom{x+y}{n} = \sum_{l=0}^n \binom{x}{l} \binom{y}{n-l} \quad (24)$$

as $x = (x_1 \cdots x_k - z)/\rho$ and $y = n-1$. Then

$$\begin{aligned} (-1)^n \frac{\widehat{c}_{n,\rho,q}^{(k)}(z)}{n!} &= \rho^n \underbrace{\int_0^1 \cdots \int_0^1}_k \sum_{m=0}^l \binom{(x_1 \cdots x_k - z)/\rho}{m} \binom{n-1}{n-m} d_q x_1 \cdots d_q x_k \\ &= \sum_{m=0}^n \binom{n-1}{m-1} \rho^n \underbrace{\int_0^1 \cdots \int_0^1}_k \binom{(x_1 \cdots x_k - z)/\rho}{m} d_q x_1 \cdots d_q x_k \\ &= \sum_{m=1}^n \binom{n-1}{m-1} \frac{c_{m,\rho,q}^{(k)}(z)}{m!}. \end{aligned}$$

Note that $\binom{n-1}{-1} = 0$. ■

Theorem 8 *For any x and y we have*

$$\begin{aligned} B_{n,\rho,q}^{(k)}(x) &= \sum_{l=0}^n \sum_{m=0}^n (-1)^{n-m} m! \rho^{n-l} S_2 \left(n, m, \frac{x}{\rho} \right) S_2 \left(m, l, \frac{y}{\rho} \right) c_{l,\rho,q}^{(k)}(y), \\ B_{n,\rho,q}^{(k)}(x) &= \sum_{l=0}^n \sum_{m=0}^n (-1)^n m! \rho^{n-l} S_2 \left(n, m, \frac{x}{\rho} \right) S_2 \left(m, l, -\frac{y}{\rho} \right) \widehat{c}_{l,\rho,q}^{(k)}(y), \\ c_{n,\rho,q}^{(k)}(x) &= \sum_{l=0}^n \sum_{m=0}^n \frac{(-1)^{n-m}}{m!} \rho^{n-l} S_1 \left(n, m, \frac{x}{\rho} \right) S_1 \left(m, l, \frac{y}{\rho} \right) B_{l,\rho,q}^{(k)}(y), \\ \widehat{c}_{n,\rho,q}^{(k)}(x) &= \sum_{l=0}^n \sum_{m=0}^n \frac{(-1)^n}{m!} \rho^{n-l} S_1 \left(n, m, -\frac{x}{\rho} \right) S_1 \left(m, l, \frac{y}{\rho} \right) B_{l,\rho,q}^{(k)}(y). \end{aligned}$$

Remark. If $\rho = 1$ and $q \rightarrow 1$, then Theorem 8 is reduced to Theorem 3.3 in [5]. If $\rho = 1$, $x = y$ and $q \rightarrow 1$, then Theorem 8 is reduced to Theorem 4.1 in [19]. A different generalization without Jackson's integrals is discussed in [18].

Proof. We shall prove the first and the fourth identities. Others can be proven similarly. By (20) in Theorem 6, and using (11), we have

$$\begin{aligned}
& \sum_{l=0}^n \sum_{m=0}^n (-1)^{n-m} m! \rho^{n-l} S_2 \left(n, m, \frac{x}{\rho} \right) S_2 \left(m, l, \frac{y}{\rho} \right) c_{l,\rho,q}^{(k)}(y) \\
&= \sum_{m=0}^n (-1)^{n-m} m! \rho^n S_2 \left(n, m, \frac{x}{\rho} \right) \sum_{l=0}^m S_2 \left(m, l, \frac{y}{\rho} \right) \rho^{-l} c_{l,\rho,q}^{(k)}(y) \\
&= \sum_{m=0}^n S_2 \left(n, m, \frac{x}{\rho} \right) \frac{(-\rho)^{n-m} m!}{[m+1]_q^k} \\
&= B_{n,\rho,q}^{(k)}(x).
\end{aligned}$$

By (19) in Theorem 6, and using (17), we have

$$\begin{aligned}
& \sum_{l=0}^n \sum_{m=0}^n \frac{(-1)^n}{m!} \rho^{n-l} S_1 \left(n, m, -\frac{x}{\rho} \right) S_1 \left(m, l, \frac{y}{\rho} \right) B_{l,\rho,q}^{(k)}(y) \\
&= \sum_{m=0}^n \frac{(-1)^n}{m!} \rho^n S_1 \left(n, m, -\frac{x}{\rho} \right) \sum_{l=0}^m S_1 \left(m, l, \frac{y}{\rho} \right) \rho^{-l} B_{l,\rho,q}^{(k)}(y) \\
&= (-1)^n \sum_{m=0}^n S_1 \left(n, m, -\frac{x}{\rho} \right) \frac{\rho^{n-m}}{[m+1]_q^k} \\
&= \widehat{c}_{n,\rho,q}^{(k)}(x).
\end{aligned}$$

■

6 Acknowledgement

This work was supported in part by the grant of Wuhan University and by the grant of Hubei Provincial Experts Program.

References

- [1] Andrews G. E., Askey R., Roy R., *Special Functions*, Encyclopedia Math. Appl., Vol.71, Cambridge Univ. Press, Cambridge, 1999.
- [2] Bayad A., Hamahata Y., *Polylogarithms and poly-Bernoulli polynomials*, Kyushu J. Math. **65** (2011), 15–24.

- [3] Carlitz L., *A note on Bernoulli and Euler polynomials of the second kind*, Scripta Math. **25** (1961), 323–330.
- [4] Carlitz L., *Weighted Stirling numbers of the first and second kind - I*, Fibonacci Quart. **18** (1980), 147–162.
- [5] Cenkci M., Komatsu T., *Poly-Bernoulli numbers and polynomials with a q parameter*, J. Number Theory, (to appear).
- [6] Comtet L., *Advanced Combinatorics*, Reidel, Dordrecht, 1974.
- [7] Coppo M.-A., Candelpergher B., *The Arakawa-Kaneko zeta functions*, Ramanujan J. **22** (2010), 153–162.
- [8] Hamahata Y., Masubuchi H., *Special multi-poly-Bernoulli numbers*, J. Integer Seq. **10** (2007), Article 07.4.1.
- [9] Jackson F. H., *On q -definite integrals*, Quart. J. **41** (1910), 193–203.
- [10] Jordan Ch., *Sur des polynômes analogues aux polynômes de Bernoulli et sur des formules de sommation analogues à celle de MacLaurin-Euler*, Acta Sci. Math. (Szeged) **4**(1928–29), 130–150.
- [11] Kamano K., Komatsu T., *Poly-Cauchy polynomials*, Mosc. J. Comb. Number Theory **3** (2013), 183–209.
- [12] Kaneko M., *Poly-Bernoulli numbers*, J. Theor. Nombres Bordeaux **9** (1997), 221–228.
- [13] Katsurada M., *Complete asymptotic expansions for certain multiple q -integrals and q -differentials of Thomae-Jackson type*, Acta Arith. **152** (2012), 109–136.
- [14] Komatsu T., *Poly-Cauchy numbers*, Kyushu J. Math. **67** (2013), 143–153.
- [15] Komatsu T., *Poly-Cauchy numbers with a q parameter*, Ramanujan J. **31** (2013), 353–371.
- [16] Komatsu T., *Sums of products of Cauchy numbers, including poly-Cauchy numbers*, J. Discrete Math. **2013** (2013), Article ID 373927, 10 pages.
- [17] Komatsu T., *Hypergeometric Cauchy numbers*, Int. J. Number Theory **9** (2013), 545–560.
- [18] Komatsu T., Liptai K., Szalay L., *Some relationships between poly-Cauchy type numbers and poly-Bernoulli type numbers*, East-West J. Math. **14** (2012), 114–120.
- [19] Komatsu T., Luca F., *Some relationships between poly-Cauchy numbers and poly-Bernoulli numbers*, Ann. Math. Inform. **41** (2013), 99–105.

- [20] Merlini D., Sprugnoli R., Verri M. C., *The Cauchy numbers*, Discrete Math. **306** (2006), 1906–1920.
- [21] Nörlund N. E., *Vorlesungen über Differenzenrechnung*, Springer, Berlin, 1924.
- [22] Riordan J., *Combinatorial identities*, John Wiley & Sons, New York, 1968